

THE ORLIK-SOLOMON ALGEBRA AND THE BERGMAN FAN OF A MATROID

ILIA ZHARKOV

ABSTRACT. Given a matroid M one can define its Orlik-Solomon algebra $OS(M)$ and the Bergman fan $\Sigma_0(M)$. On the other hand to any rational polyhedral fan Σ one can associate its tropical homology and cohomology groups $\mathcal{F}_\bullet(\Sigma)$, $\mathcal{F}^\bullet(\Sigma)$. We will show that the projective Orlik-Solomon algebra $OS_0(M)$ is canonically isomorphic to $\mathcal{F}^\bullet(\Sigma_0(M))$.

This note was meant to serve as an appendix to the tropical homology paper [IKMZ12]. But since a shorter geometric argument will appear in the realizable case (all that is needed there) in [IKMZ12] the algebraic proof given here makes now more sense as an independent article.

1. NOTATIONS AND STATEMENTS

1.1. Tropical homology and cohomology. Let $\Sigma = \bigcup \sigma \subset \mathbb{R}^N = \mathbb{Z}^N \otimes \mathbb{R}$ be an integral polyhedral fan. For each cone $\sigma \subset \Sigma$ we denote by $\langle \sigma \rangle_{\mathbb{Z}}$ the integral lattice in the vector subspace linearly spanned by σ .

Definition 1. [IKMZ12] The *homology* group $\mathcal{F}_k(\Sigma)$ is the subgroup of $\wedge^k \mathbb{Z}^N$ generated by the elements $v_1 \wedge \cdots \wedge v_k$, where all $v_1, \dots, v_k \in \langle \sigma \rangle_{\mathbb{Z}}$ for some cone $\sigma \in \Sigma$. The *cohomology* is the dual group $\mathcal{F}^k(\Sigma) := \text{Hom}(\mathcal{F}_k(\Sigma), \mathbb{Z})$, which is the quotient of $\wedge^\bullet(\mathbb{Z}^N)^*$ by $(\mathcal{F}_k)^\perp$.

Lemma 2. *The wedge product on $\wedge^\bullet(\mathbb{Z}^N)^*$ descends to \mathcal{F}^\bullet , that is, \mathcal{F}^\bullet is endowed with a natural algebra structure over \mathbb{Z} .*

Proof. We just need to show that the subgroup of $\wedge^\bullet(\mathbb{Z}^N)^*$ annihilating \mathcal{F}_\bullet forms an ideal. Let $f \in (\mathcal{F}_k)^\perp$, then for any $\alpha \in (\mathbb{Z}^N)^*$ and any collection $v_0, v_1, \dots, v_k \in \langle \sigma \rangle_{\mathbb{Z}}$ we have

$$(\alpha \wedge f)(v_0 \wedge v_1 \wedge \cdots \wedge v_k) = \sum_{i=0}^k (-1)^i \alpha(v_i) f(v_0 \wedge \dots \wedge \hat{v}_i \wedge \dots \wedge v_k),$$

The research is partially supported by the NSF FRG grant DMS-0854989.

which vanishes since any k -subset of v_0, v_1, \dots, v_k is also in $\langle \sigma \rangle_{\mathbb{Z}}$. Hence $\alpha \wedge f$ is in $(\mathcal{F}_{k+1})^\perp$. \square

1.2. The Bergman fan. Let M be a loopless matroid of rank n on the set $\{0, \dots, N\}$. Let V be the rank $N + 1$ free abelian group generated by elements e_0, \dots, e_N . Consider the simplicial fan $\Sigma(M) \subset V_{\mathbb{R}}$ built on the lattice of flats of M . Namely, the rays of Σ are along the vectors $e_J := e_{j_1} + \dots + e_{j_k}$ for each flat $J = \{j_1, \dots, j_k\}$. The k dimensional cones of $\Sigma(M)$ are spanned by the k -tuples of rays indexed by flags of flats of length k . We will also use the notation

$$E_I := e_{i_1} \wedge \dots \wedge e_{i_k}$$

for any subset $\{i_1, \dots, i_k\} \subset M$. Note the distinction between E_I and e_I . We reserve letter J to denote flats in M , while I will be used for general subsets of M .

The Bergman fan is the quotient fan $\Sigma_0(M)$ of $\Sigma(M)$ at the ray e_M . Namely, it is defined like above by the lattice of proper flats of M in the quotient lattice $V_0 = V / \langle e_0 + \dots + e_N \rangle$.

Sturmfels [St02] noticed the importance of the Bergman fan to tropical geometry where it represents a linear space. Later Ardila and Klivans [AK06] studied its combinatorics and showed, among other things, that $\Sigma_0(M)$ is indeed a balanced fan of degree 1. Here we adopted the definition from [KP11].

1.3. The Orlik-Solomon algebra. To the same matroid M one can associate its Orlik-Solomon algebra $OS^\bullet(M)$ over \mathbb{Z} defined below. In case M is realizable by a hyperplane arrangement in \mathbb{P}^{n-1} , the projectivized version $OS_0^\bullet(M)$ of this algebra calculates the cohomology of the complement of this arrangement. (See [OT92] for more details).

Let W be the rank $N + 1$ free abelian group generated by elements f_0, \dots, f_N . Then $OS^\bullet(M) := \wedge^\bullet W / \mathcal{I}^\bullet$, where the Orlik-Solomon ideal \mathcal{I} is generated by the elements

$$\partial(f_{i_0} \wedge f_{i_1} \wedge \dots \wedge f_{i_k}) := \sum_{s=0}^k (-1)^s f_{i_0} \wedge \dots \wedge \hat{f}_{i_s} \wedge \dots \wedge f_{i_k},$$

for all dependent subsets $I = \{i_0, i_1, \dots, i_k\}$. We will use the notation

$$F_I := f_{i_0} \wedge f_{i_1} \wedge \dots \wedge f_{i_k}.$$

The sign of F_I depends on the order of I , so we assume that all subsets of M are ordered.

The projective Orlik-Solomon algebra $OS_0^\bullet(M)$ is defined as follows. Let W_0 be the subgroup of W generated by all differences $f_i - f_j$.

Then we set $OS_0^\bullet(M) := \wedge^\bullet W_0 / \mathcal{I}_0^\bullet$, where \mathcal{I}_0 is the restriction of \mathcal{I} to the subalgebra $\wedge^\bullet W_0 \subset \wedge^\bullet W$.

Theorem 3. *Consider V and W above as dual groups with the dual bases $\{e_0, \dots, e_N\}$ and $\{f_0, \dots, f_N\}$. Then $\mathcal{F}_k(\Sigma(M))^\perp = \mathcal{I}^k$.*

An important corollary of this theorem is our main result:

Theorem 4. *There is a canonical isomorphism $\mathcal{F}^\bullet(\Sigma_0(M)) \cong OS_0^\bullet(M)$ of graded algebras.*

2. PROOF OF THEOREMS 3 AND 4

For a flat $J \subset M$ we consider the restricted groups $\mathcal{F}_k(J) := \mathcal{F}_k(\Sigma(J))$ as subgroups of $\wedge^\bullet V$ under the natural embedding $\wedge^\bullet(\mathbb{Z}\langle e_j, j \in J \rangle) \subset \wedge^\bullet V$. We also consider the restricted Orlik-Solomon algebra $OS^\bullet(J) := \wedge^\bullet W / \mathcal{I}^\bullet(J)$ by defining the ideal $\mathcal{I}^\bullet(J) \subset \wedge^\bullet W$ to be generated by the ∂F_I with dependent $I \subset J$, and by the $f_i, i \notin J$.

Lemma 5. $\mathcal{F}_k(M) = \mathbb{Z}\langle \mathcal{F}_k(J) \rangle_{\text{rk}(J)=k}$.

Proof. Let $J \subset J''$ be two flats whose ranks differ by 2 or more. Let J'_1, \dots, J'_s be the set of flats between J and J'' of rank exactly one larger than the rank of J . Then the sets $J, J'_1 \setminus J, \dots, J'_s \setminus J$ give a partition of J'' . Hence

$$e_J \wedge e_{J''} = \sum_{i=1}^s e_J \wedge e_{J'_i}.$$

By induction, for any k -flag of flats $J_1 \subset \dots \subset J_k$ the element $e_{J_1} \wedge \dots \wedge e_{J_k}$ can be rewritten as a sum

$$e_{J_1} \wedge \dots \wedge e_{J_k} = \sum e_{J''_1} \wedge \dots \wedge e_{J''_k},$$

where all flags $J''_1 \subset \dots \subset J''_k$ consist of flats of ranks $1, \dots, k$, respectively. \square

Lemma 6. *As an abelian group $\mathcal{I}^k = \mathbb{Z}\langle F_{I'}, \partial F_{I''} \rangle$, where I' and I'' run over dependent sets in M of size k and $k+1$, respectively. In particular, in the top degree $\mathcal{I}^n = \mathbb{Z}\langle F_I, \text{Im}\{\partial : \wedge^{n+1} W \rightarrow \wedge^n W\} \rangle = \mathbb{Z}\langle F_I, \ker\{\partial : \wedge^n W \rightarrow \wedge^{n-1} W\} \rangle$, where I runs over dependent sets of size n .*

Proof. The statement follows from the Leibnitz rule:

$$\partial F_{I'} \wedge F_{I''} = \partial(F_{I'} \wedge F_{I''}) \pm F_{I'} \wedge \partial F_{I''}.$$

For the top degree note that every $(n+1)$ -set is dependent and $(\wedge^\bullet W, \partial)$ is an acyclic complex, that is $\text{Im } \partial = \ker \partial$. \square

Remark. For the second subset of generators it is enough to take ∂F_I with rank of I exactly k , since ∂F_I with I of smaller ranks are already included in the first subset of generators.

Remark. Note that the projective Orlik-Solomon ideal \mathcal{I}_0 is generated as an abelian group just by the ∂F_I for dependent I . Indeed, note that $\wedge^\bullet W_0 = \text{Im}\{\partial : \wedge^\bullet W \rightarrow \wedge^\bullet W\} = \ker\{\partial : \wedge^\bullet W \rightarrow \wedge^\bullet W\}$. But from the Lemma 6 if $\alpha \in \mathcal{I}$ we can write $\alpha = \sum F_{I'} + \sum \partial F_{I''}$, with all I', I'' dependent sets. On the other hand $\partial\alpha = \partial(\sum F_{I'}) = 0$ means $\sum F_{I'} = \partial \sum F_{\hat{I}}$, where every \hat{I} is an extension by one element of some dependent I , and hence is also dependent.

Lemma 7. $\mathcal{I}^k = \cap_{\text{rk}(J)=k} \mathcal{I}^k(J)$.

Proof. First we argue that for any rank k flat J we have $\mathcal{I}^k \subset \mathcal{I}^k(J)$. Indeed, according to Lemma 6 and the first remark after it we just have to show that $\partial F_{\hat{I}} \in \mathcal{I}^k(J)$ for \hat{I} of size $k+1$ and rank k . If $\hat{I} \subset J$ or $|\hat{I} \setminus J| \geq 2$, we are done. Otherwise, say $|\hat{I} \setminus J| = \{s\}$. Then

$$\partial F_{\hat{I}} = f_s \wedge (\dots) \pm F_{\hat{I} \setminus s}.$$

But $\hat{I} \setminus s \subset J$ must have rank $k-1$ (or, otherwise $\hat{I} \subset J$), hence is dependent. Thus $F_{\hat{I} \setminus s}$ is in $\mathcal{I}^k(J)$, and so is $\partial F_{\hat{I}}$. Consequently, $\mathcal{I}^k \subset \cap_{\text{rk}(J)=k} \mathcal{I}^k(J)$.

To show the converse we notice that each I is contained in a unique flat of the same rank (the matroidal closure of I). We group the terms in an element $\alpha = \sum F_I \in \wedge^k W$ by their flats:

$$\alpha = \alpha_{<k} + \sum_{\text{rk}(J')=k} \alpha_{J'}$$

where $\alpha_{<k}$ contains terms F_I with dependent I .

Now if $\alpha \in \mathcal{I}^k(J)$ for some rank k flat J , then in the above decomposition $\alpha_{<k} \in \mathcal{I}^k \subset \mathcal{I}^k(J)$. Also all $\alpha_{J'}$ with $J' \neq J$, are in $\mathcal{I}^k(J)$, and hence so is α_J . But all terms F_I in $\mathcal{I}^k(J)$ with independent $I \subset J$ have to come from $\partial F_{\hat{I}}$ for some dependent $\hat{I} \subset J$. Thus $\alpha_J \in \mathcal{I}^k$. Taking the intersection over all k -flats completes the proof. \square

Proof of Theorem 3. Taking the intersection in Lemma 7 is orthogonal to taking the sum in Lemma 5. Thus it is enough to prove the statement in the top degree for any matroid. By dualizing the top degree part of Lemma 6 it suffices then to show that $\mathcal{F}_n = \langle E_I \rangle \cap \text{Im}(\partial^*)$, where the I run over independent n -sets of M and $\partial^* : \wedge^{n-1} V \rightarrow \wedge^n V$, $\partial^*(E_I) = e_M \wedge E_I$ is the adjoint operator to $\partial : \wedge^n W \rightarrow \wedge^{n-1} W$.

For any complete flag of flats $J_1 \subset \cdots \subset J_{n-1} \subset M$ the polyvector

$$e_{J_1} \wedge \cdots \wedge e_M = e_{J_1} \wedge e_{J_2 \setminus J_1} \wedge \cdots \wedge e_{M \setminus J_{n-1}} = \sum E_I$$

contains only terms with independent I . Thus $\mathcal{F}_n \subset \langle E_I \rangle \cap \text{Im}(\partial^*)$. We will show the converse by induction on the rank of M . For rank 1 matroids both spaces are $\mathbb{Z}\langle e_M \rangle$ and there is nothing to prove.

Suppose now $\alpha = e_M \wedge \beta = \sum E_{\hat{I}}$, with all \hat{I} independent. We may choose a representation for $\beta = \sum E_I$ with all I independent subsets as follows. Substituting, say, $e_0 = -\sum_{i=1}^N e_i \bmod e_M$ into β we will have

$$\alpha = e_0 \wedge \beta + (\text{terms with no } e_0)$$

and any term E_I with dependent I in β will result in $E_{0 \cup I}$ in α with dependent $\hat{I} = 0 \cup I$ which cannot happen.

Let J_1, \dots, J_r be all rank $(n-1)$ flats in M . We again group the terms in β by the respective flats $\beta = \beta_{J_1} + \cdots + \beta_{J_r}$. Then writing

$$\alpha = (e_{J_1} \wedge \beta_{J_1} + \cdots + e_{J_r} \wedge \beta_{J_r}) + (e_{M \setminus J_1} \wedge \beta_{J_1} + \cdots + e_{M \setminus J_r} \wedge \beta_{J_r})$$

we note that both α and the second summand contain terms E_I only with independent I . On the other hand, each $e_{J_k} \wedge \beta_{J_k}$ contains terms of rank $n-1$, and there no cancellations possible among different k . Thus $e_{J_k} \wedge \beta_{J_k} = 0$. By exactness of the $(e_{J_k} \wedge)$ -operator we can write each β_{J_k} as $e_{J_k} \wedge \gamma_{J_k}$ and use the induction assumption. \square

Proof of Theorem 4. With the choice of the dual bases for V and W the restriction to W_0 in W is exactly dual to the quotient by e_M in V , and the duality extends to the exterior algebras. On the other hand, the identification in Theorem 3 clearly extends to the level of graded ideals $\mathcal{F}_\bullet(\Sigma(M))^\perp = \mathcal{I}^\bullet$, as well as to their restrictions to $\wedge^\bullet W_0$. \square

Acknowledgments. As I already mentioned this note was intended to be an appendix to our (long overdue) joint project on tropical homology [IKMZ12] with Ilia Itenberg, Ludmil Katzarkov and Grisha Mikhalkin. Of course, numerous discussions with all three of them were crucial for the formulation and the proof of the main statement. I am very grateful to my coauthors for their permission to publish the result as a separate article. I would also like to thank Federico Ardila for several very helpful conversations. Finally I should mention that another inductive proof of Theorem 4 was recently given by Kristin Shaw in her thesis [Sh12] using tropical modifications.

REFERENCES

- [AK06] Federico Ardila and Caroline Klivans. *The Bergman Complex of a Matroid and Phylogenetic Trees*. J. Comb. Theory, Ser. B 96(1): 38-49 (2006).
- [IKMZ12] Ilia Itenberg, Ludmil Katzarkov, Grigory Mikhalkin and Ilia Zharkov. *Tropical Homology*. In preparation.
- [KP11] Eric Katz and Sam Payne. *Realization spaces for tropical fans*. Combinatorial aspects of commutative algebra and algebraic geometry, Abel Symp., 6, Springer 2011, 73–88.
- [OT92] Peter Orlik and Hiroaki Terao. *Arrangements of hyperplanes*. Springer Verlag, 1992.
- [Sh12] Kristin Shaw. *Solving systems of polynomial equations*. Available at <http://www.math.toronto.edu/shawkm/>
- [St02] Bernd Sturmfels. *Solving systems of polynomial equations*. CBMS Regional Conference Series in Mathematics, vol. 97. American Mathematical Society, Providence, RI, 2002.

KANSAS STATE UNIVERSITY, 138 CARDWELL HALL, MANHATTAN, KS 66506
E-mail address: zharkov@math.ksu.edu